

## Roberto Caldeyro-Barcia (1921?1996) <sup>[1]</sup>

By: Mandile, Olivia Keywords: Obstetrics <sup>[2]</sup>

Roberto Caldeyro-Barcia studied fetal health in Uruguay during the second half of the twentieth century. Caldeyro-Barcia developed Montevideo units, which are used to quantify intrauterine pressure, or the force of contractions during labor. Intrauterine pressure is a useful measure of the progression of labor and the health of a [fetus](#) <sup>[3]</sup>. Caldeyro-Barcia's research on fetal health often contradicted common obstetric practices, prompting him to publically challenge practices such as [induction](#) <sup>[4]</sup> of labor using oxytocin, forced pushing during labor, and birth position in which the woman lays on her back during labor. Caldeyro-Barcia's methods of monitoring intrauterine pressure and development of Montevideo units furthered research in maternal and fetal health and improved the use of medical interventions during labor and delivery.

Caldeyro-Barcia was born on 26 September 1921 in Montevideo, Uruguay. His mother, Elvira Barcia, came from a family with a medical background and his father, Joaquin Caldeyro, was a physician. Caldeyro-Barcia attended primary school at the English School in Montevideo, where he met fellow student Ofelia Stajano, whom he would later marry. As a teenager, Caldeyro-Barcia attended the Elbro Fernandez School in Montevideo, and began to court Stajano.

In 1938, Caldeyro-Barcia entered the University of Uruguay's Faculty of Medicine in Montevideo, where he studied medicine. During his time in school, Caldeyro-Barcia and Stajano became engaged. The couple married a year later in January 1946 and had six children, Ofelia, Lucia, Roberto, Carlos, Eduardo, and Martin.

At the University of Uruguay, Caldeyro-Barcia researched contractions of the [uterus](#) <sup>[5]</sup> during labor alongside Hermógenes Alvarez, an associate professor in obstetrics and gynecology. During delivery, a pregnant woman's [uterus](#) <sup>[5]</sup> contracts to push the [fetus](#) <sup>[3]</sup> out. During their research together, Caldeyro-Barcia and Alvarez obtained the first known recording of intrauterine pressure, or the force of those contractions. To measure intrauterine pressure, they placed a microballoon in the [uterus](#) <sup>[5]</sup>. As the [uterus](#) <sup>[5]</sup> contracted, the balloon was compressed. That compression indicated the pressure of the contraction. Using the data gathered through microballoons, Caldeyro-Barcia and Alvarez then compared the force of contractions during normal labors and labors that were not progressing normally.

In 1947, Caldeyro-Barcia graduated with honors from the University of Uruguay in Montevideo, Uruguay, with a medical degree. After graduation, he became an assistant professor of physiology at the University of Uruguay. He continued to work with Alvarez, and shortly after obtaining the first recording of intrauterine pressure, they developed Montevideo units to quantify the force of contractions. Montevideo units are calculated by subtracting the baseline pressure of a [uterus](#) <sup>[5]</sup> from the highest pressure of each contraction throughout ten minutes, then summing the pressures. They are used during delivery to help physicians monitor the progress of labor. If the sum of the pressures is approximately two hundred

Montevideo units, then labor is progressing as needed. However, if the force of contractions are significantly below the normal range, the contractions are not strong enough to dilate, or open, the [cervix](#) [6] wide enough for the [fetus](#) [3] to enter the birth canal. The invention of Montevideo units by Caldeyro-Barcia and Alvarez was an important development for physicians to monitor labor. According to Otto Armas, professor in gynecology at the Central University of Venezuela in Caracas, Venezuela, the invention of Montevideo units resulted in more rational and calculated use of oxytocin, a drug commonly used to progress labor.

Throughout the next decade, Caldeyro-Barcia and Alvarez studied the effects of uterine contractions on fetal heart rate. In the 1950s, Caldeyro-Barcia and Alvarez found that during contractions the fetal heart rate goes down. They noted two different types of decreased fetal heart rate, Type I dips and Type II dips. In Type I dips, the fetal heart rate falls during the contraction as a result of head compression and recovers immediately after the contraction is over. In Type II dips, the fetal heart rate falls during the contraction but takes a significant amount of time to recover once the contraction is over. Type II dips can indicate that the [fetus](#) [3] is not receiving enough oxygen, which can result in adverse health effects and calls for medical intervention to deliver the infant safely. In 1952, Caldeyro-Barcia and Alvarez presented their research on monitoring intrauterine pressure and fetal heart rate during labor in England. Following their presentation, their techniques for monitoring contraction force and fetal heart rate became widely accepted within the medical community.

In 1959, University of Uruguay appointed Caldeyro-Barcia chairman of the newly created Department of Obstetric Physiology. During his time as chairman, he began studying the effects of oxytocin on the [uterus](#) [5]. Physicians routinely administered oxytocin, a [hormone](#) [7] released during contractions and labor, to artificially induce labor in pregnant women. Though [induction](#) [4] by oxytocin was common at the time, the side effects of the procedure were unknown. To study potential side effects, Caldeyro-Barcia monitored fetal health during oxytocin induced deliveries. He found that thirty-eight percent of oxytocin induced contractions caused Type II dips, which indicated that the fetus's health was compromised. Caldeyro-Barcia also found that Type II dips caused by oxytocin induced contractions may even result in fetal death.

In 1969, Caldeyro-Barcia organized a conference titled "Perinatal Factors Affecting Human Development" to address obstetric interventions such as [induction](#) [4] by oxytocin. At that conference, he presented his research on oxytocin and Type II dips, and argued that oxytocin should only be used when medically necessary and not in inductions that are elective, or not medically necessitated.

In the 1960s, Caldeyro-Barcia researched labor activity, including methods for delaying preterm labor. Caldeyro-Barcia and his colleagues began developing tocolytics, medications used to halt [premature labor](#) [8] by inhibiting contractions of the [uterus](#) [5]. According to Caldeyro-Barcia's colleague Erich Staling, prior to tocolytic medications, preterm labor would often result in Cesarean sections, a procedure in which the [fetus](#) [3] is surgically removed from the woman's [uterus](#) [5]. Cesarean sections often resulted in higher rates of maternal and infant complications. The development of tocolytics enabled physicians to prevent many preterm births and reduce the frequency of invasive measures like Cesarean sections.

In 1970, Caldeyro Barcia became the first director of the Latin American Center of Perinatology and Human Development in Montevideo. During his tenure as the director, Caldeyro-Barcia explained labor as a natural activity and stressed that the role of a physician

or midwife was to facilitate the process rather than to intervene. To preserve the natural physiology of labor, he created a birthing chair that allowed laboring mothers to remain in an upright position, as opposed to the more common position where laboring women lay on their backs. In the upright position that Caldeyro-Barcia advocated for, a pregnant woman's pelvic opening is wider than it is while lying down, allowing labor to progress more quickly. A faster labor also decreases the need for forceps or Cesarean sections. The upright position also lessens the pain of contractions by working with gravity instead of against it, pushing the fetus [3] into the birth canal. Several years later, Caldeyro-Barcia presented his recommendations on maternal posture during childbirth at a symposium in Rio de Janeiro, Brazil. The *West of England Medical Journal* published those recommendations posthumously.

In 1977, Caldeyro-Barcia studied the effects of forced pushing during labor on fetal health. At the time, physicians and midwives encouraged laboring mothers to push forcefully and for a prolonged amounts of time to speed delivery. Caldeyro Barcia's research demonstrated that forced pushing during labor results in an insufficient oxygen supply to the fetus [3]. Insufficient oxygen supply can damage the fetus [3]. For that reason, Caldeyro-Barcia encouraged laboring mothers to push when they naturally felt the need, rather than when their physician encouraged it.

At the age of 60, Caldeyro-Barcia retired from his position as director of the Latin American Center of Perinatology and Human Development. He then became the Director of the Maternal-Child Health Program at the Ministry of Public Health of Uruguay in Montevideo. Throughout his career, Caldeyro-Barcia travelled frequently, and while travelling he found that many Uruguayan scientists had left Uruguay for countries with better working conditions. In 1987, Caldeyro-Barcia helped establish the Basic Sciences Development Program in Uruguay in Montevideo, Uruguay. The program was created to facilitate the return to Uruguay of scientists who had left the country and who wanted to return.

Caldeyro-Barcia served as president of numerous organizations, including the Department of Obstetric Physiology at the University of Uruguay, the Latin American Center of Perinatology and Human Development in Montevideo, Uruguay, the International Federation of Gynecologists and Obstetricians in London, United Kingdom, and the Basic Sciences Development Program in Montevideo, Uruguay. He also cofounded the *Journal of Perinatal Medicine*, an international journal that covers topics in the field of obstetrics and neonatology, including pregnancy [9], labor, and the time immediately following birth. Caldeyro-Barcia was nominated three times for a Nobel Prize, although he never received the award.

Toward the end of his life, Caldeyro-Barcia developed diabetes and heart disease. He died at the age of seventy-five in Montevideo on 2 November 1996, following heart surgery.

## Sources

1. Armas, Otto Rodriguez. "In Memoriam: Professor Roberto Caldeyro-Barcia." *International Journal of Gynecology and Obstetrics* 59 (1997): 175?6.
2. Caldeyro-Barcia, Roberto, and Juan J. Poseiro. "Physiology of the uterine contraction." *Clinical Obstetrics and Gynecology* 3 (1960): 386?410.
3. Caldeyro-Barcia, Roberto, and Peter. M. Dunn. "Appropriate maternal posture during childbirth." *Parto Humanizado: Formas Alternativas* (Delivery Made Human: Alternative

- Options). Campinas, Brazil: Publ: Editora de Unicamp (1992): 17?25.
4. Capurro, Haroldo, and José Belizán. "Roberto Caldeyro-Barcia: An Appreciation." *Paediatric and Perinatal Epidemiology* 12 (1998): 240?2.
  5. Crowther, Damien. GPNotebook. ?Cardiotocographic dips.?
  6. Dunn, Peter M. "Perinatal Profiles: Roberto Caldeyro-Barcia: Obstetric Physiologist Extraordinary." *NeoReviews* 9 (2008): 187?91.
  7. Haire, Doris. "In Memory of Roberto Caldeyro-Barcia, MD, 1921?1996." *Birth* 24 (1997): 211?2.
  8. Haas, David, Tara Benjamin, Sawyer Renata, and Sara Quinney. ?Short-term tocolytics for preterm delivery?Current Perspectives.? *International Journal of Women?s Health* 6 (2014): 343?349. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3971910/> <sup>[10]</sup> (Accessed June 12, 2017).
  9. McCormack, Patricia. ?Birthing chair shortens second stage of labor.? UPI. (1981).
  10. Noble, Elizabeth. "A Tribute to Robert Caldeyro-Barcia, MD." *Birth* 24 (1997): 212?3.
  11. Pose, S. V., J. B. Castillo, E. O. Mora-Rojas, A. Soto-Yances, and Roberto Caldeyro-Barcia. "Test of fetal tolerance to induced uterine contractions for the diagnosis of chronic distress." *Perinatal Factors Affecting Human Development* 96 (1969): 96?104
  12. Saling, Erich. ?Obituary.? *Journal of Perinatal Medicine* 25 (1997): 9?10
  13. Whalley, Eric, and Alan J. Riley. "Effect of an intra-uterine micro-balloon on the duration of parturition in the *rat* <sup>[11]</sup>." *Journal of Endocrinology* 72 (1977): 411?2.

Roberto Caldeyro-Barcia studied fetal health in Uruguay during the second half of the twentieth century. Caldeyro-Barcia developed Montevideo units, which are used to quantify intrauterine pressure, or the force of contractions during labor. Intrauterine pressure is a useful measure of the progression of labor and the health of a fetus. Caldeyro-Barcia's research on fetal health often contradicted common obstetric practices, prompting him to publically challenge practices such as induction of labor using oxytocin, forced pushing during labor, and birth position in which the woman lays on her back during labor. Caldeyro-Barcia's methods of monitoring intrauterine pressure and development of Montevideo units furthered research in maternal and fetal health and improved the use of medical interventions during labor and delivery.

## Subject

Programa para el Desarrollo de las Ciencias Básicas (Uruguay) <sup>[12]</sup> Programme for the Development of the Basic Sciences (Uruguay) <sup>[13]</sup> PEDECIBA <sup>[14]</sup> Fetal assessment <sup>[15]</sup> Fetal surveillance <sup>[16]</sup> Fetus--Monitoring <sup>[17]</sup> Monitoring, Fetal <sup>[18]</sup> Childbirth <sup>[19]</sup> Delivery (Obstetrics) <sup>[20]</sup> Fetal heart rate monitoring <sup>[21]</sup> Obstetrics <sup>[22]</sup> Obstetric Labor, Premature <sup>[23]</sup> Obstetric Labor Complications <sup>[24]</sup> OBSTETRIC LABOR <sup>[25]</sup> Pregnancy <sup>[26]</sup> Gynecology <sup>[27]</sup> Tocolytic Agents <sup>[28]</sup> Tocolysis <sup>[29]</sup> Heart Rate, Fetal <sup>[30]</sup> Uterine Contraction <sup>[31]</sup> Cardiotocography <sup>[32]</sup> Oxytocin <sup>[33]</sup>

## Topic

People <sup>[34]</sup>

## Publisher

Arizona State University. School of Life Sciences. Center for Biology and Society. Embryo

Project Encyclopedia.

## Rights

Copyright Arizona Board of Regents Licensed as Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported (CC BY-NC-SA 3.0)  
<http://creativecommons.org/licenses/by-nc-sa/3.0/>

## Format

Articles <sup>[35]</sup>

## Last Modified

Wednesday, July 4, 2018 - 04:40

## DC Date

2017-07-02

## DC Date Accessioned

Sunday, July 2, 2017 - 22:15

## DC Date Available

Sunday, July 2, 2017 - 22:15

## DC Date Created

2017-07-02

## DC Date Created Standard

Sunday, July 2, 2017 - 07:00

- [Contact Us](#)

© 2018 Arizona Board of Regents

- The Embryo Project at Arizona State University, 1711 South Rural Road, Tempe  
Arizona 85287, United States

---

**Source URL:** <https://embryo.asu.edu/pages/roberto-caldeyro-barcia-1921-1996>

### Links:

- [1] <https://embryo.asu.edu/pages/roberto-caldeyro-barcia-1921-1996>
- [2] <https://embryo.asu.edu/keywords/obstetrics>
- [3] <https://embryo.asu.edu/search?text=fetus>
- [4] <https://embryo.asu.edu/search?text=induction>
- [5] <https://embryo.asu.edu/search?text=uterus>

- [6] <https://embryo.asu.edu/search?text=cervix>
- [7] <https://embryo.asu.edu/search?text=hormone>
- [8] <https://embryo.asu.edu/search?text=premature%20labor>
- [9] <https://embryo.asu.edu/search?text=pregnancy>
- [10] <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3971910/>
- [11] <https://embryo.asu.edu/search?text=rat>
- [12] <https://embryo.asu.edu/library-congress-subject-headings/programa-para-el-desarrollo-de-las-ciencias-basicas-uruguay>
- [13] <https://embryo.asu.edu/library-congress-subject-headings/programme-development-basic-sciences-uruguay>
- [14] <https://embryo.asu.edu/library-congress-subject-headings/pedeciba>
- [15] <https://embryo.asu.edu/library-congress-subject-headings/fetal-assessment>
- [16] <https://embryo.asu.edu/library-congress-subject-headings/fetal-surveillance>
- [17] <https://embryo.asu.edu/library-congress-subject-headings/fetus-monitoring>
- [18] <https://embryo.asu.edu/library-congress-subject-headings/monitoring-fetal>
- [19] <https://embryo.asu.edu/library-congress-subject-headings/childbirth>
- [20] <https://embryo.asu.edu/library-congress-subject-headings/delivery-obstetrics>
- [21] <https://embryo.asu.edu/library-congress-subject-headings/fetal-heart-rate-monitoring>
- [22] <https://embryo.asu.edu/medical-subject-headings/obstetrics>
- [23] <https://embryo.asu.edu/medical-subject-headings/obstetric-labor-premature>
- [24] <https://embryo.asu.edu/medical-subject-headings/obstetric-labor-complications>
- [25] <https://embryo.asu.edu/medical-subject-headings/obstetric-labor>
- [26] <https://embryo.asu.edu/medical-subject-headings/pregnancy>
- [27] <https://embryo.asu.edu/medical-subject-headings/gynecology>
- [28] <https://embryo.asu.edu/medical-subject-headings/tocolytic-agents>
- [29] <https://embryo.asu.edu/medical-subject-headings/tocolysis>
- [30] <https://embryo.asu.edu/medical-subject-headings/heart-rate-fetal>
- [31] <https://embryo.asu.edu/medical-subject-headings/uterine-contraction>
- [32] <https://embryo.asu.edu/medical-subject-headings/cardiocography>
- [33] <https://embryo.asu.edu/medical-subject-headings/oxytocin>
- [34] <https://embryo.asu.edu/topics/people>
- [35] <https://embryo.asu.edu/formats/articles>