

## Morphogenesis <sup>[1]</sup>

By: Sunderland, Mary E. Keywords: Morphogenesis <sup>[2]</sup> Differentiation <sup>[3]</sup>

The term morphogenesis generally refers to the processes by which order is created in the developing organism. This order is achieved as differentiated cells carefully organize into tissues, organs, organ systems, and ultimately the organism as a whole. Questions centered on morphogenesis have aimed to uncover the mechanisms responsible for this [organization](#) <sup>[5]</sup>, and developmental biology textbooks have identified morphogenesis as one of the main challenges in the field. The concept of morphogenesis is intertwined with those of [differentiation](#) <sup>[6]</sup>, growth, and reproduction. Each comprises the fundamental components of development that have commonly been used to categorize the problems that motivate developmental biology.

Previously, morphogenesis had appeared in discussions of [embryology](#) <sup>[7]</sup>, particularly those regarding the processes responsible for the form of the organism. After the discovery of the [organizer](#) <sup>[8]</sup> by [Hans Spemann](#) <sup>[9]</sup> and [Hilde Mangold](#) <sup>[10]</sup> in 1927 there was a widespread effort to study those regions of the embryo that were capable of inducing the formation of body structures. This search for the nature of the [organizer](#) <sup>[8]</sup> also led many to adopt a biochemical approach to [embryology](#) <sup>[7]</sup> in attempts to characterize the inductor substances and understand their action.

[Paul Weiss](#) <sup>[11]</sup> used a sculptor analogy to describe morphogenesis in his 1939 book *Principles of Development*. He distinguished morphogenesis from growth and explained that growth was the creation of mass whereas morphogenesis was the shaping of that mass. This process of morphogenesis was exemplified by amphibian [gastrulation](#) <sup>[12]</sup>, which had been meticulously described by [Johannes Holtfreter](#) <sup>[13]</sup> and also by asexual reproduction in [Volvox](#) <sup>[14]</sup>.

In 1934 [Julian Huxley](#) <sup>[15]</sup> and [Gavin De Beer](#) <sup>[16]</sup> published *Elements of Experimental Embryology* <sup>[17]</sup> wherein they discussed Spemann's [organizer](#) <sup>[8]</sup> in the context of Charles Manning Child's [gradient theory](#) <sup>[18]</sup>. Together, they suggested that these two interpretations offered an excellent theoretical framework to think about development, particularly problems of morphogenesis. Spemann, however, was not convinced by Child's theory, which he found to be an over-simplified explanation, and therefore was also highly critical of Huxley and De Beer's contribution.

As evidence about the processes of morphogenesis accumulated additional concepts were created. For example, morphogenetic field and morphogenetic substratum were terms used by [Joseph Needham](#) <sup>[19]</sup> in his embryological text *Biochemistry and Morphogenesis* <sup>[4]</sup> (1950). Needham defined morphogenesis as the process by which an organism acquired its characteristic form.

In 1952 [John Tyler Bonner](#) <sup>[20]</sup> published *Morphogenesis: An Essay on Development* <sup>[21]</sup> wherein he surveyed a diverse selection of developmental phenomena across a wide range of organisms to determine if he could find shared processes. Development was described as

being composed of both constructive and limiting forces; those that built up the organism and those that kept it in check. Bonner identified the constructive processes as growth, [differentiation](#) [6], and morphogenetic movements?those processes that involved cellular movement. Bonner, like Weiss, described them as analogous to the process of a sculptor shaping clay. Bonner suggested that cellular slime molds offered an alternative approach to studying morphogenesis as their life cycle separated periods of growth and [differentiation](#) [6] from those of morphogenesis.

In the 1970s morphogenesis was still identified as one of the least understood aspects of development and was described as the progressive acquisition of form during development. A number of systems and processes were studied to investigate this problem such as limb development in chicks, limb regeneration in [amphibians](#) [22], and the development of [neural crest cells](#) [23] .

Today those who study morphogenesis are asking many questions and trying to determine, for example, how tissues form from populations of cells, how tissues construct organs, how organs grow, how growth is coordinated, how migrating cells are oriented, and how [polarity](#) [24] is achieved. The problem of morphogenesis is recognized by many to be one of the most elusive questions of development as it is intertwined with questions of [regulation](#) [25] and how the organism functions as a whole.