

Gastrulation in *Xenopus* ^[1]

By: Maayan, Inbar Keywords: Amphibians ^[2] Frogs ^[3] Gastrulation ^[4]

The process of [gastrulation](#) ^[6] allows for the formation of the [germ layers](#) ^[7] in metazoan embryos, and is generally achieved through a series of complex and coordinated cellular movements. The process of [gastrulation](#) ^[6] can be either diploblastic or triploblastic. In diploblastic organisms like cnidaria or ctenophora, only the [endoderm](#) ^[8] and the [ectoderm](#) ^[9] form; in triploblastic organisms (most other complex metazoans), triploblastic [gastrulation](#) ^[6] produces all three [germ layers](#) ^[7]. The [gastrula](#) ^[10], the product of [gastrulation](#) ^[6], was named by [Ernst Haeckel](#) ^[11] in the mid-1870s; the name comes from Latin, where *gaster* means stomach, and indeed the gut (archenteron) is one of the most distinctive features of the [gastrula](#) ^[10].

Since the early twentieth century, experimental embryologists like [Hans Spemann](#) ^[12] and [Wilhelm Roux](#) ^[13] have extensively studied [gastrulation](#) ^[6] in amphibian embryos in an attempt to learn more about how establishment of different regions in the body is determined. The size and structure of [Xenopus laevis](#) ^[14] (African clawed frogs) embryos have made the species into a [model organism](#) ^[15] for early developmental study. The following is a detailed explanation of [gastrulation](#) ^[6] in *Xenopus*; while [gastrulation](#) ^[6] varies across species, studies in *Xenopus* have shed considerable light on the process in general.

The raw material for [gastrulation](#) ^[6] is the [blastula](#) ^[16], a hollow sphere of cells; the space inside of the [blastula](#) ^[16] is called the [blastocoel](#) ^[17]. If the [blastula](#) ^[16] were compared to a globe of the world, the North Pole would correspond to the animal pole of the embryo, and the South Pole to the embryo's vegetal pole. The hemispheres of the [blastula](#) ^[16] correspond to the names of their respective poles. Cellular division occurs much more rapidly near the animal (active) pole of the [frog](#) ^[18] embryo than near the yolkier vegetal (sedentary) pole; the [yolk](#) ^[19] provides the embryo with nutrition, but slows down cell division around it. The equator between the two hemispheres is called the marginal zone. The [blastocoel](#) ^[17] occupies most of the inside of the animal hemisphere, since the cells in the vegetal pole bear the [yolk](#) ^[19] of the embryo and therefore occupy more of the inner space in that region.

While the animal-vegetal gradient is determined in the [egg](#) ^[20] prior to [fertilization](#) ^[21], the point of entry of the [sperm](#) ^[22] lends the [frog](#) ^[18] its dorsal-ventral (back-and-front) axis. The prospective ventral side of the embryo is on the side of the sperm's entry, while the prospective dorsal side, the side at which the [blastopore](#) ^[23] forms, is opposite the sperm's point of entry. The [blastopore](#) ^[23] is a groove in the side of the embryo that results from the invagination (the formation of a groove) of a small group of future endodermal cells, and forms right below the equator of the embryo, in the marginal zone. Here, cells undergo apical constriction to become "bottle" cells (tapered cells resembling narrow Erlenmeyer flasks); since most of the cytoplasm in each of these cells is made to migrate toward the center of the embryo, the part of the cell in contact with the groove to become much narrower. The lip of the [blastopore](#) ^[23] nearer to the animal pole is termed the dorsal lip of the [blastopore](#) ^[23].

Once the [blastopore](#) [23] has formed, the cells of the animal hemisphere undergo epiboly (flattening of the outer layer of cells and radial intercalation of the layer of cells beneath them) and move toward the [blastopore](#) [23]. As these cells divide and extend to envelope the inside of the developing [frog](#) [18], the marginal zone cells involute (or fold inward) at the dorsal lip of the [blastopore](#) [23]. During involution, these inner cells move toward the dorsal lip and then fold over themselves and progress toward the animal pole (the top of the embryo). While the location at which the involuting cells change direction remains the dorsal lip of the [blastopore](#) [23], the cells passing through that point change: as the original tapered bottle cells involute, they are replaced by precursor cells for the head [mesoderm](#) [24], followed by chordamesoderm cells. The latter will go on to form the [notochord](#) [25] of the embryo. It is important to note that this involution depends on the movement of the deep marginal zone cells, as opposed to the movement of the superficial bottle cells of the [blastopore](#) [23].

While cells are involuting and the pre-[ectoderm](#) [9] is going through epiboly, the [blastocoel](#) [17] moves ventrally (up and away from the dorsal lip of the [blastopore](#) [23], and then eventually around toward the vegetal pole) and shrinks as the involuting cells progressively move in to occupy the region, much like how frost creeps up a window on a cold day. The involuting marginal zone cells progress ventrally to line the roof of the archenteron (the gut) which expands in proportion to the disappearance of the [blastocoel](#) [17]. As it forms, the archenteron itself resembles a balloon that is being inflated through the [blastopore](#) [23] and into the animal hemisphere.

When epiboly and involution near completion, the material of the vegetal hemisphere closest to the [blastopore](#) [23] becomes constricted by the dorsal and ventral [blastopore](#) [23] lips to form a circular [yolk](#) [19] plug. This [yolk](#) [19] plug then continues to shrink as ectodermal epiboly progresses, and is eventually absorbed into the embryo; the inner layer, formed from vegetal pole cells, becomes the [endoderm](#) [8]. The [mesoderm](#) [24] constitutes the region between the superficial [ectoderm](#) [9] and the internal [endoderm](#) [8], and lines the upper part of the archenteron. At the end of [gastrulation](#) [6], the *Xenopus* embryo becomes known as a [gastrula](#) [10]. It has an [ectoderm](#) [9] as its outermost layer, and an [endoderm](#) [8] as its innermost layer. The [mesoderm](#) [24] lies between the two layers, and the [endoderm](#) [8] and [mesoderm](#) [24] line the archenteron of the [gastrula](#) [10].

While *Xenopus* serves as the [model organism](#) [15] for study of amphibian [gastrulation](#) [6], some key differences exist in the process across the class. The amphibian class is made up of three major orders: anura (tailless [amphibians](#) [26] like frogs), caudata/ urodela (tailed [amphibians](#) [26] like salamanders) and caecilians (legless [amphibians](#) [26], [snake](#) [27]-like). Whereas in *Xenopus* the pre-mesodermal tissues originate exclusively from the deeper marginal zone, urodeles derive mesodermal tissue from both superficial and deeper marginal zone cells. In addition, in the 1940s German embryologist [Johannes Holtfreter](#) [28] was able to show that the dorsal marginal zone cells, i.e., the bottle cells of the [blastopore](#) [23], serve a crucial role in involution and can even induce a [blastopore](#) [23] to form in urodeles. This is not the case with *Xenopus*, which indeed relies on the bottle cells for initiation, but not for the continuance of the [gastrulation](#) [6].

Gastrulation is crucial to the proper development of all multicellular animals, especially those with complex tissue structure. Designation of the [ectoderm](#) [9] is essential for the later formation of the epidermis (skin) and nervous system. The [mesoderm](#) [24] gives rise to the skeleton, muscles, connective tissue, blood and internal organs. The [endoderm](#) [8], the

innermost germ layer, goes on to form other organs like the lungs, liver, and pancreas as well as the lining of the digestive tract.

Studies of [gastrulation](#) [6] in *Xenopus* have allowed developmental biologists to construct accurate [fate maps](#) [29] and to make inferences regarding the [ontogeny](#) [30] of [amphibians](#) [26]. Amphibians have therefore provided the means for a better understanding of the process by which a mass of relatively undifferentiated cells transforms into an intricate and functional organism. While *Xenopus*, in particular, entered the field of [embryology](#) [31] as an already-established clinical tool for effective [pregnancy](#) [32] assays, this [frog](#) [18] species soon became a prominent [model organism](#) [15] for experimental embryologists, and continues to aid greatly in the investigation of [ontogeny](#) [30].

Sources

1. Beetschen, Jean-Claude. "Amphibian Gastrulation: History and Evolution of a 125 Year-Old Concept," *International Journal of [Developmental Biology](#)* [33] 45 (2001): 771-95.
2. Dale, Leslie. "Vertebrate Development: Multiple Phases to Endoderm Formation," *Current Biology* 9 (1999): R812-15.
3. Gilbert, Scott F. *[Developmental Biology](#)* [33], 4th ed. Sunderland: Sinauer, 1994.
4. Gurdon, John B., and [Nick Hopwood](#) [34]. "The Introduction of *Xenopus laevis* [14] into [Developmental Biology](#) [33]: Of Empire, Pregnancy Testing and Ribosomal Genes," *International Journal of [Developmental Biology](#)* [33] 44 (2000): 43-50.
5. Stern, Claudio D., ed. *Gastrulation: From Cells to Embryo*. Cold Spring Harbor: [Cold Spring Harbor Laboratory](#) [35] Press, 2004.

The process of gastrulation allows for the formation of the germ layers in metazoan embryos, and is generally achieved through a series of complex and coordinated cellular movements. The process of gastrulation can be either diploblastic or triploblastic. In diploblastic organisms like cnidaria or ctenophora, only the endoderm and the ectoderm form; in triploblastic organisms (most other complex metazoans), triploblastic gastrulation produces all three germ layers. The gastrula, the product of gastrulation, was named by Ernst Haeckel in the mid-1870s; the name comes from Latin, where gaster means stomach, and indeed the gut (archenteron) is one of the most distinctive features of the gastrula.

Subject

Haeckel, Ernst, 1834-1919 [36] Gastrulation [37]

Topic

Processes [38]

Publisher

Arizona State University. School of Life Sciences. Center for Biology and Society. Embryo Project Encyclopedia.

Rights

© Arizona Board of Regents Licensed as Creative Commons Attribution-NonCommercial-Share Alike 3.0 Unported (CC BY-NC-SA 3.0) <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Format

Articles ^[39]

Last Modified

Wednesday, July 4, 2018 - 04:40

DC Date Accessioned

Thursday, May 10, 2012 - 14:06

DC Date Available

Thursday, May 10, 2012 - 14:06

DC Date Created

2011-03-10

DC Date Created Standard

Thursday, March 10, 2011 - 07:00

- [Contact Us](#)

© 2018 Arizona Board of Regents

- The Embryo Project at Arizona State University, 1711 South Rural Road, Tempe
Arizona 85287, United States

Source URL: <https://embryo.asu.edu/pages/gastrulation-xenopus>

Links:

- [1] <https://embryo.asu.edu/pages/gastrulation-xenopus>
- [2] <https://embryo.asu.edu/keywords/amphibians>
- [3] <https://embryo.asu.edu/keywords/frogs>
- [4] <https://embryo.asu.edu/keywords/gastrulation>
- [5] <https://embryo.asu.edu/search?text=Gastrulation%20in%20Xenopus>
- [6] <https://embryo.asu.edu/search?text=gastrulation>
- [7] <https://embryo.asu.edu/search?text=germ%20layers>
- [8] <https://embryo.asu.edu/search?text=endoderm>
- [9] <https://embryo.asu.edu/search?text=ectoderm>
- [10] <https://embryo.asu.edu/search?text=gastrula>
- [11] <https://embryo.asu.edu/search?text=Ernst%20Haeckel>

- [12] <https://embryo.asu.edu/search?text=Hans%20Spemann>
- [13] <https://embryo.asu.edu/search?text=Wilhelm%20Roux>
- [14] <https://embryo.asu.edu/search?text=Xenopus%20laevis>
- [15] <https://embryo.asu.edu/search?text=model%20organism>
- [16] <https://embryo.asu.edu/search?text=blastula>
- [17] <https://embryo.asu.edu/search?text=blastocoel>
- [18] <https://embryo.asu.edu/search?text=frog>
- [19] <https://embryo.asu.edu/search?text=yolk>
- [20] <https://embryo.asu.edu/search?text=egg>
- [21] <https://embryo.asu.edu/search?text=fertilization>
- [22] <https://embryo.asu.edu/search?text=sperm>
- [23] <https://embryo.asu.edu/search?text=blastopore>
- [24] <https://embryo.asu.edu/search?text=mesoderm>
- [25] <https://embryo.asu.edu/search?text=notochord>
- [26] <https://embryo.asu.edu/search?text=amphibians>
- [27] <https://embryo.asu.edu/search?text=snake>
- [28] <https://embryo.asu.edu/search?text=Johannes%20Holtfreter>
- [29] <https://embryo.asu.edu/search?text=fate%20maps>
- [30] <https://embryo.asu.edu/search?text=ontogeny>
- [31] <https://embryo.asu.edu/search?text=embryology>
- [32] <https://embryo.asu.edu/search?text=pregnancy>
- [33] <https://embryo.asu.edu/search?text=Developmental%20Biology>
- [34] <https://embryo.asu.edu/search?text=Nick%20Hopwood>
- [35] <https://embryo.asu.edu/search?text=Cold%20Spring%20Harbor%20Laboratory>
- [36] <https://embryo.asu.edu/library-congress-subject-headings/haeckel-ernst-1834-1919>
- [37] <https://embryo.asu.edu/medical-subject-headings/gastrulation>
- [38] <https://embryo.asu.edu/topics/processes>
- [39] <https://embryo.asu.edu/formats/articles>